

# Ragweed: diffusional spread and pollen load

**Robert Richter<sup>1\*</sup>, Michael Leitner<sup>2</sup>, Gero Vogl<sup>1</sup>**

<sup>1</sup> Fakultät für Physik, Universität Wien, Vienna, Austria

<sup>2</sup> Heinz Maier-Leibnitz Zentrum (MLZ), Technische Universität München, Munich, Germany  
[\\*robert.richter@univie.ac.at](mailto:robert.richter@univie.ac.at)

Invasive alien species cause a major threat to our ecosystems and health [1]. We simulate the spread of *Ambrosia artemisiifolia* L. (common ragweed) under current climate as well as with climate change assuming 1.5°C temperature increase from the year 1990 until 2050 (Fig. 1). *Ambrosia artemisiifolia* L. spreading rapidly in Europe [2-5] is an annual weed that causes pollen allergies [6]. Therefore, it is necessary to implement pollen load into the spread model to be able to determine the pollen pressure in the study area Austria and Bavaria. Recently, this was done for whole Europe on a coarse scale of 50 x 50 km with two different models [7]. We, in contrast, used a finer scale of about 6 x 6 km and considered in addition a higher number of pollen traps in relation to the size of the study area. For calculating pollen dispersal, we assumed that the drifting time for each pollen grain (i.e. the time it stays in the air) follows an exponential distribution with parameter  $\tau$  and the empirical distribution of wind directions and velocities is constant for each individual trajectory. To achieve an estimate of the spatial pollen distribution, we compared the annual pollen totals recorded at eight Austrian pollen traps with the results of our simulations (Fig. 2) and thereby determined  $\tau$  to be 1.72 h [8].




Figure 1: Distribution of grid cells infested by *Ambrosia artemisiifolia* L. in 2005 (black squares, left) and in 2050 (right, with climate change, different colours indicate the probability of infestation).




Figure 2: Simulated pollen dispersion (darkest shade indicates maximum) and location of eight Austrian pollen traps (blacksquares with numbers).

## References

- [1] F. Essl, S. Dullinger, I. Kleinbauer: *Changes in the spatio-temporal patterns and habitat preferences of Ambrosia artemisiifolia during its invasion of Austria*. Preslia, **81**(2), 119–133 (2009)
- [2] M. Smolik, S. Dullinger, F. Essl, I. Kleinbauer, M. Leitner, J. Peterseil, L.-M. Stadler, G. Vogl: *Integrating species distribution models and interacting particle systems to predict the spread of an invasive alien plant*. J. Biogeogr., **37**, 411–422 (2010); R. Richter, S. Dullinger, F. Essl, M. Leitner, G. Vogl: *How to account for habitat suitability in weed management programmes?* Biol. Invasions, **15**, 657–669 (2013)
- [3] S. Cunze, M.C. Leiblein, O. Tackenberg: *Range expansion of Ambrosia artemisiifolia in Europe is promoted by climate change*, SRN Ecology [www.hindawi.com/isrn/ecology/2013/610126/](http://www.hindawi.com/isrn/ecology/2013/610126/) (2013)
- [4] F. Essl et al: *Biological Flora of the British Isles: Ambrosia artemisiifolia*. Journal of Ecology, doi: 10.1111/1365-2745.12424 (2015)
- [5] D. Chapman, T. Haynes, S. Beal, F. Essl, J. Bullock: *Phenology predicts the native and invasive range limits of common ragweed*. Global Change Biology **20**, 192–202 (2014)
- [6] M. Smith, L. Cecchi, C. Skjøth, G. Karrer, B. Šikoparija.: *Common ragweed: A threat to environmental health in Europe*. Environment International, **61**, 115–126 (2013)
- [7] L. Hamaoui-Laguel et al: *Effects of climate change and seed dispersal on airborne ragweed pollen loads in Europe*. Nature Climate Change, doi: 10.1038/NCLIMATE2652 (2015)
- [8] R. Richter, U.E. Berger, S. Dullinger, F. Essl, M. Leitner, M. Smith, G. Vogl: *Spread of invasive ragweed: climate change, management and how to reduce allergy costs*. Journal of Applied Ecology, **50**(6), 1422–1430 (2013); R. Richter: *Dissertation*, Universität Wien (2015)